



TRYPTIC SOY AGAR

- For in vitro use only -

Plated Media

PT80 – Tryptic Soy Agar (TSA)
PT81 – TSA (SXT)
PT89 – TSA w Yeast Extract
PB75 – TSA w 5% Sheep Blood
PB81 – TSA w 7% Sheep Blood
PB69 – TSA w 5% Horse Blood
PB80 – TSA w 7% Horse Blood

Tryptic Soy Agar (TSA) is a general purpose plating medium used for the isolation, cultivation, and maintenance of a variety of fastidious and non-fastidious microorganisms.

Leavitt et al. demonstrated the versatility of TSA by cultivating both aerobic and anaerobic microbes using TSA. TSA is recognized and recommended by numerous agencies around the world. Our standard formulation is prepared according to the United States Pharmacopeia (USP) and recommended for various different applications put forth by the Association of Official Analytical Chemists (AOAC), the International Dairy Federation (IDF), the United States Department of Agriculture (USDA), and the American Public Health Association (APHA).

Tryptic Soy Agar is a highly nutritious base that meets the growth requirements of many types of microorganisms including bacteria, yeasts, and molds. Many modifications have been to the TSA formulation to increase both its nutritious and selective properties. TSA with 5% defibrinated sheep blood is used extensively for the cultivation and recovery of fastidious microbial species and for the determination of hemolytic reactions that are important differential characteristic especially among the streptococci. This medium is also suitable for performing the overnight CAMP test; Group B streptococci produce an extracellular substance (CAMP factor) that can act synergistically with the beta-toxin produced by some *Staphylococcus aureus* strains. When placed in close proximity to one another the two organisms produce a zone of increased hemolysis.

Tubed Media

TT80 – TSA Slant
TT80-18 – TSA Pour Plate [18-mL]
TB75 – TSA Blood Slant

The CAMP test can also be used to help identify pathogenic species of *Listeria*.

TSA with horse blood is used to isolate more fastidious organisms. Horse blood contains both X and V factor, which are essential growth factors for some organisms such as *Haemophilus* species. Sheep and human blood are not suitable since they contain specific enzymes that inactivate V Factor.

Although, some laboratories prefer a plated medium with a higher blood content (7-10%) or with horse blood, these mediums should not be used for determination of hemolytic reactions or for the CAMP test. The increased blood content can make hemolytic reactions less distinct and more difficult to read while defibrinated horse blood, in some instances, has shown to give hemolytic reactions different from sheep blood.

TSA supplemented with yeast extract is described in the FDA Bacteriological Analytical Manual for the isolation and purification of *Listeria monocytogenes* as well as other heterotrophic organisms. Gunn, Ohashi, Gaydos, and Holt developed the selective SXT formulation by adding the selective agents sulfamethoxazole and trimethoprim. They found that this medium gave superior isolation of group A and B streptococci from throat specimens by inhibiting the growth of the normal throat flora.

Plain TSA can also be used in the differentiation of *Haemophilus* species when used in conjunction with X, V, and XV factor disks. Differentiation is based on the growth pattern around the various disks.

Formulation per Litre of Medium

PT80 & TT80 Tryptic Soy Agar

Pancreatic Digest of Casein.....	15.0 g
Papaic Digest of Soybean Meal.....	5.0 g
Sodium Chloride	5.0 g
Agar	15.0 g

pH 7.3 ± 0.2

Additional Ingredients per Liter:

PB75 & TB75 TSA with 5% Sheep Blood

Defibrinated Sheep Blood.....	50.0 mL
-------------------------------	---------

PB69 TSB with 5% Horse Blood

Defibrinated Horse Blood.....	50.0 mL
-------------------------------	---------

PB80 TSA with 7% Horse Blood

Defibrinated Horse Blood.....	70.0 mL
-------------------------------	---------

PB81 TSA with 7% Sheep Blood

Defibrinated Sheep Blood.....	70.0 mL
-------------------------------	---------

PT81 TSA (SXT)

Defibrinated Sheep Blood.....	50.0 mL
Sulfamethoxazole	23.75 µg
Trimethoprim	1.25 µg

PT89 TSA with Yeast Extract

Yeast Extract	6.0 g
---------------------	-------

Recommended Procedure

(Please refer to appropriate literature for a more detailed procedure)

1. Allow medium to adjust to room temperature prior to inoculation.
2. Inoculate by performing a four-quadrant streak on the plated media to obtain well-isolated colonies. For tubed media, streak the surface of the medium in a fishtail motion from bottom up.
3. For TSA with blood, several stabs should be made into the medium during inoculation to better detect beta-hemolysis reactions.
4. Incubate aerobically or in CO₂-rich

environment at 35°C (plates should be inverted).

5. Examine plates and tubes after 18 to 24 hours and at 48 hours.

CAMP Procedure

1. Allow medium to adjust to room temperature and ensure that the plate surface is dry prior to inoculation.
2. Obtain a pure overnight culture of *Staphylococcus aureus* ATCC 25923 or 33862. With an inoculating needle or edge of a loop streak *Staphylococcus* in a straight line across the center of the plate.
3. Streak test organism in a straight line 2 to 3cm long and perpendicular to the staphylococci streak. The line should come close (approx 3mm) but not touch the staphylococci streak. Four test streaks can be performed on each plate although one of the streaks should be a known positive (*Streptococcus agalactiae* ATCC 12386).
4. Label the streaks on the bottom of the plate (media side).
5. Incubate plates aerobically in an inverted position at 35°C.
6. Examine plates after 18 to 24 hours.

Interpretation of Results

TSA with 5% Defibrinated Sheep Blood is commonly used as a primary plating medium. Primary isolation is performed to separate and isolate organisms present in a specimen. This separation allows for characterization of colony types and may indicate the presence of clinically significant bacteria. When examining primary plates a hand lens or stereoscopic microscope should be available for examining very small colonies. The different types of colonial morphology appearing on the agar plate should be noted as well as the number of each morphotype present. Hemolysis is a useful differential characteristic that is best viewed when a bright light is transmitted from behind the plate. Four different types of hemolysis can be described:

1. Alpha-hemolysis (α) – Partial hemolysis that results in a greenish discoloration around the colony
2. Beta-hemolysis (β) – Complete lysis of red blood cells resulting in a clear zone around the colony
3. Gamma-hemolysis (γ) – No hemolysis resulting in no change in the medium
4. Alpha-prime-hemolysis (α') – A small zone of complete hydrolysis that is surrounded by an area of partial hemolysis

Additional results such as pigment production and odor should also be recorded.

The CAMP test can be used to presumptively identify group B streptococci (*S. agalactiae*). A positive CAMP reaction is defined by the production of a distinct arrowhead zone of complete hemolysis at the point of intersection between the test streak and the *S. aureus* streak. The hemolysis reaction must extend throughout the depth of the agar plate. A negative CAMP reaction is no arrowhead phenomenon or a slight increased zone of hemolysis but with no arrowhead formation. Some organisms such as group A streptococci may show increased hemolysis at the zone of intersection.

Additional tests should be performed on isolated colonies from pure culture in order to complete identification.

- *For TSA with sheep blood, stabbing into the medium during inoculation creates an area of reduced oxygen tension that is necessary for hemolysis by oxygen-labile hemolysin O*
- *Before performing the CAMP test, a beta-hemolytic organism must be presumptively identified as a member of the genus Streptococcus by a catalase test and gram stain*
- *Bacteria, other than group B streptococci, may give a positive CAMP reaction, such as Pasteurella haemolytica, Listeria monocytogenes, Burkholderia pseudomallei, Corynebacterium renale, Mobiluncus mulieris, Mobiluncus curtisii, and Propionibacterium*

Quality Control

After checking for correct pH, colour, depth, and sterility, the following organisms are used to determine the growth performance of the completed medium.

Organism	Expected Results
TSA	
<i>Pseudomonas aeruginosa</i> ATCC 27853	Growth
<i>Streptococcus pneumoniae</i> ATCC 6305	Growth

TSA w/ 5% Sheep Blood

<i>Streptococcus pneumoniae</i> ATCC 6305	Growth, α -hemolysis
<i>Streptococcus pyogenes</i> ATCC 19615	Growth, β -hemolysis, CAMP (–)
<i>Streptococcus agalactiae</i> ATCC 12386	Growth, β -hemolysis, CAMP (+)
<i>Escherichia coli</i> ATCC 25922	Growth

TSA (SXT)

<i>Streptococcus agalactiae</i> ATCC 12386	Growth
<i>Streptococcus pyogenes</i> ATCC 19615	Growth
<i>Streptococcus pneumoniae</i> ATCC 6305	Inhibition
<i>Escherichia coli</i> ATCC 25922	Inhibition

TSA w/ Yeast Extract

<i>Listeria monocytogenes</i> ATCC 19114	Growth
---	--------

TSA w/ Horse Blood

<i>Haemophilus influenzae</i> ATCC 10211	Growth
<i>Haemophilus haemolyticus</i> ATCC 33390	Growth

Storage and Shelf Life

Our various Tryptic Soy Agar formulations should be stored away from direct light at 4°C to 8°C. For plated media, the medium side should be uppermost to prevent excessive accumulation of moisture on the agar surface. Under these conditions this mediums have the following shelf lives from the date of manufacture:

- PT80 – TSA – 12 weeks
- PT81 – TSA (SXT) – 8 weeks
- PT89 – TSA w Yeast Extract – 12 weeks
- PB75 – TSA w 5% Sheep Blood – 8 weeks
- PB81 – TSA w 7% Sheep Blood – 8 weeks
- PB69 – TSA w 5% Horse Blood – 8 weeks
- PB80 – TSA w 7% Horse Blood – 8 weeks
- TT80 – TSA Slant – 16 weeks
- TB75 – TSA Blood Slant – 8 weeks

References

1. Leavitt JM, Naidorf IJ, Shugaevsky P. The undetected anaerobe in endodontics; a sensitive medium for detection of both aerobes and anaerobes. NY J Dentist 1955; 25:377-82.
2. Fraser G. A plate method for the rapid identification of *Listeria (Erysipelothrix) monocytogenes*. Vet Rec 1962; 74:50-1.
3. Darling CL. Standardization and evaluation of CAMP reaction for the prompt, presumptive identification of *Streptococcus agalactiae* (Lancefield group B). J Clin Micro 1975; 1:171-4.
4. Gunn BA, Ohashi DK, Gaydos CA, Holt ES. Selective and enhanced recovery of group A and B streptococci from throat cultures with sheep blood containing sulfamethoxazole and trimethoprim. J Clin Micro 1977; 5:650-5.
5. Kurzynski TA, Meise CK. Evaluation of sulfamethoxazole-trimethoprim blood agar plates for recovery of group A streptococci from throat cultures. J Clin Micro 1979; 9:189.
6. MacFaddin JF. Media for isolation-cultivation-maintenance of medical bacteria, Vol I. Baltimore: Williams & Wilkins, 1985.
7. Isenberg HD, Ed. Clinical microbiology procedures handbook. Washington, DC: ASM, 1992.
8. Orth DS. Handbook of cosmetic microbiology. New York: Marcel Dekker Inc., 1993.
9. Greenberg AE, Clesceri LS, Eaton AD, Eds. Standard methods for the examination of water and wastewater. 19th ed. Washington, DC: APHA, 1995.
10. USP. The United States pharmacopeia. 23rd ed. Rockville, MD: United States Pharmacopeial Convention, 1995.
11. NCCLS. Quality assurance for commercially prepared microbiological culture media. 2nd ed. NCCLS document M22-A2. Wayne, PA: NCCLS, 1996.
12. Murray PR, Baron EJ, Pfaller MA, Tenover FC, Tenover RH. Manual of clinical microbiology. 7th ed. Washington: ASM, 1999.
13. FDA. Bacteriological analytical manual, 2002. Retrieved January 27, 2003, from FDA website: <http://www.cfsan.fda.gov>.

Original: January 2003

Revised / Reviewed: October 2014